

がん化学療法レジメン

食道癌・胆道癌・肛門管癌・膵癌

疾患名	療法名	催吐性	制吐剤	投与量	投与日	薬剤名	投与量	投与日	投与期間・1クール
食道癌	FP(70-700)+RT ※RTは月～金 1.8-2Gy/Fr/Day(計50.4～60Gy)	H	プロイメンド	150mg	day1	CDDP	70mg/m <sup>2</sup>	day1	4週間
			アロキシ	0.75mg		5-FU	700mg/m <sup>2</sup>	day1-4	
			DEX注	9.9-6.6-13.2mg	day1-3				
	FP(80-800)	H	プロイメンド	150mg	day1	CDDP	80mg/m <sup>2</sup>	day1	4週間
			アロキシ	0.75mg		5-FU	800mg/m <sup>2</sup>	day1-5	
			DEX注	9.9-6.6-13.2mg	day1-3				
	NAC FP(80-800)	H	プロイメンド	150mg	day1	CDDP	80mg/m <sup>2</sup>	day1	3週間
			アロキシ	0.75mg		5-FU	800mg/m <sup>2</sup>	day1-5	
DEX注			9.9-6.6-13.2mg	day1-3					
Neda/5FU(90-800)±RT	M	アロキシ	0.75mg	day1	Neda	90mg/m <sup>2</sup> ※高齢者では 80mg/m <sup>2</sup> を推奨	day1	4週間 RTは5週間	
		DEX注 DEX錠	6.6mg 8mg	day1 day2-3	5-FU	800mg/m <sup>2</sup>	day1-5		
Weekly PTX(100)	L	DEX注	9.9mg	day1,8,15, 22,29,36	PTX	100mg/m <sup>2</sup>	day1,8,15,22,29,36	7週間	
DOC	L	DEX注	6.6mg	day1	DOC	70mg/m <sup>2</sup>	day1	3週間	
Nivolumab	Mini					Nivo	240mg/body	day1	2週間
mFOLFOX6 入院/SV2.5使用	M	アロキシ	0.75mg	day1	L-OHP	85mg/m <sup>2</sup>	day1	2週間	
		DEX注 DEX錠	9.9mg 8mg	day1 day2-3	I-LV	200mg/m <sup>2</sup>			
						5-FU	ポース 400mg/m <sup>2</sup> 持注 2,400mg/m <sup>2</sup> (入院は1,200mg/m <sup>2</sup> ×2回)		day1 day1-2 (46hr)
胆道癌	GCS StageIV	H	アプレピタント	125-80-80mg	day1-3	CDDP	25mg/m <sup>2</sup>	day1	2週間 ※12コース
			アロキシ	0.75mg	day1	GEM	1,000mg/m <sup>2</sup>	day1	
			DEX注 DEX錠	9.9mg 8mg	day1 day2-3	S1	80mg/m <sup>2</sup> /日 ※規格 25mg/20mg	day1-7	
	GEM+CDDP	M	アロキシ	0.75mg	day1,8	GEM	1,000mg/m <sup>2</sup>	day1,8	3週間
			DEX注 DEX錠	9.9mg 8mg	day1,8 day2-3 day9-10	CDDP	25mg/m <sup>2</sup>		
GEM(2投1休)	L	DEX注	6.6mg	day1,8	GEM	1,000mg/m <sup>2</sup>	day1,8	3週間	
GEM(3投1休)	L	DEX注	6.6mg	day1,8,15	GEM	1,000mg/m <sup>2</sup>	day1,8,15	4週間	
肛門管癌	5-FU+MMC+RT	L	DEX注	6.6mg	day1,29	MMC	10mg/m <sup>2</sup>	day1,29	6週間
							5-FU	1,000mg/m <sup>2</sup>	
膵癌	FOLFIRINOX 入院 SV2.5使用	H	アプレピタント	125-80-80mg	day1-3	L-OHP	85mg/m <sup>2</sup>	day1	2週間
			アロキシ	0.75mg	day1	I-LV	200mg/m <sup>2</sup>		
			DEX注 DEX錠	9.9mg 8mg	day1 day2-4	CPT-11	180mg/m <sup>2</sup>		
							5-FU	ポース 400mg/m <sup>2</sup> 持注 2,400mg/m <sup>2</sup> (入院は1,200mg/m <sup>2</sup> ×2回)	
	GEM+nab-PTX	M	アロキシ	0.75mg	day1,8,15	GEM	1,000mg/m <sup>2</sup>	day1,8,15	4週間
			DEX注	9.9mg	day1,8,15	nab-PTX	125mg/m <sup>2</sup>		
			DEX錠	8mg	day2-3, 9-10, 16-17				
	NAC GS 切除可能な膵癌	M	DEX注	6.6mg	day1,8	GEM	1,000mg/m <sup>2</sup>	day1,8	3週間 ※2コース
				S1	80mg/m <sup>2</sup> /日 ※規格 25mg/20mg	day1-14			
nal-CPT-11+I-LV/5-FU 入院/SV2.5使用	M	アロキシ	0.75mg	day1	nal- CPT-11	70mg/m <sup>2</sup>	day1	2週間	
		DEX注 DEX錠	9.9mg 8mg	day1 day2-3	I-LV	200mg/m <sup>2</sup>	day1		
						5-FU	持注 2,400mg/m <sup>2</sup> (入院は1,200mg/m <sup>2</sup> ×2回)		day1 day1-2 (46hr)
GEM(2投1休)	L	DEX注	3.3mg	day1,8	GEM	1,000mg/m <sup>2</sup>	day1,8	3週間	
GEM(3投1休)	L	DEX注	3.3mg	day1,8,15	GEM	1,000mg/m <sup>2</sup>	day1,8,15	4週間	